

3.36pt



Mathematical  
Institute

# The Nazarov-Sodin constant and critical points of Gaussian fields

M. McAULEY

*Joint work with Dmitry Beliaev and Stephen Muirhead*  
Mathematical Institute  
University of Oxford

Random Waves in Oxford, June 2018

Oxford  
Mathematics

Oxford  
Mathematics



Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be a stationary Gaussian field with zero-mean, unit variance and covariance function  $\kappa : \mathbb{R}^2 \rightarrow [-1, 1]$  and spectral measure  $\rho$ , i.e. for  $x, y \in \mathbb{R}^2$

$$\kappa(x) = \mathbb{E}(f(y)f(y+x)) = \int_{\mathbb{R}^2} e^{it \cdot x} d\rho(t)$$

Basic assumptions:

1.  $\kappa \in C^{4+}(\mathbb{R}^2)$  (which implies  $f \in C^{2+}(\mathbb{R}^2)$  a.s.)
2.  $\nabla^2 f(0)$  is a non-degenerate Gaussian vector

We are interested in the geometry of the level sets

$$\{f = \ell\} := \{x \in \mathbb{R}^2 \mid f(x) = \ell\}$$

for  $\ell \in \mathbb{R}$ .

For  $\Omega \subset \mathbb{R}^2$  let  $N_{LS}(\ell, \Omega)$  be the number of components of  $\{f = \ell\}$  in  $\Omega$ .

## Theorem (Nazarov-Sodin 2016)

If  $f$  is ergodic then there exists  $c_{NS}(\rho) \geq 0$  such that

$$N_{LS}(0, R \cdot \Omega) / (\text{Area}(\Omega) R^2) \rightarrow c_{NS}(\rho)$$

a.s. and in  $L^1$ .

## Theorem (Kurlberg-Wigman 2018)

If  $\rho$  has compact support then there exists  $c_{NS}(\rho) \geq 0$  such that

$$\mathbb{E}(N_{LS}(0, [0, R]^2)) = c_{NS}(\rho) R^2 + O(R)$$

Moreover  $c_{NS}(\rho)$  is continuous in  $\rho$  (w.r.t. the  $w^*$ -topology).

For  $\Omega \subset \mathbb{R}^2$  let  $N_{LS}(\ell, \Omega)$  be the number of components of  $\{f = \ell\}$  in  $\Omega$ .

**Theorem (Nazarov-Sodin 2016)**

*If  $f$  is ergodic then there exists  $c_{NS}(\rho, \ell) \geq 0$  such that*

$$N_{LS}(\ell, R \cdot \Omega) / (\text{Area}(\Omega) R^2) \rightarrow c_{NS}(\rho, \ell)$$

*a.s. and in  $L^1$ .*

**Theorem (Kurlberg-Wigman 2018)**

*If  $\rho$  has compact support then there exists  $c_{NS}(\rho, \ell) \geq 0$  such that*

$$\mathbb{E}(N_{LS}(\ell, [0, R]^2)) = c_{NS}(\rho, \ell) R^2 + O(R)$$

*Moreover  $c_{NS}(\rho, \ell)$  is continuous in  $\rho$  (w.r.t. the  $w^*$ -topology) for each  $\ell \in \mathbb{R}$ .*

For  $\Omega \subset \mathbb{R}^2$  let  $N_{ES}(\ell, \Omega)$  be the number of components of  $\{f \geq \ell\}$  in  $\Omega$ .

Theorem (Nazarov-Sodin 2016)

If  $f$  is ergodic then there exists  $c_{ES}(\rho, \ell) \geq 0$  such that

$$N_{ES}(\ell, R \cdot \Omega) / (\text{Area}(\Omega) R^2) \rightarrow c_{ES}(\rho, \ell)$$


a.s. and in  $L^1$ .

Theorem (Kurlberg-Wigman 2018)

If  $\rho$  has compact support then there exists  $c_{ES}(\rho, \ell) \geq 0$  such that

$$\mathbb{E}(N_{ES}(\ell, [0, R]^2)) = c_{ES}(\rho, \ell) R^2 + O(R)$$

Moreover  $c_{ES}(\rho, \ell)$  is continuous in  $\rho$  (w.r.t. the  $w^*$ -topology) for each  $\ell \in \mathbb{R}$ .



$$\#\{\text{Components of } \{f = l\}\} \approx \#\{\text{Components of } \{f \geq l\}\} \\ + \#\{\text{Components of } \{f \leq l\}\}$$


## Corollary

$$c_{NS}(\rho, l) = c_{ES}(\rho, l) + c_{ES}(\rho, -l)$$

## Definition

If  $f$  is aperiodic we say that a saddle point  $x$  is *lower connected* if it is in the closure of only one component of  $\{f < l\}$ . We say that  $x$  is *upper connected* if it is in the closure of only one component of  $\{f > l\}$ .

(When  $f$  is periodic, we use a different definition for lower/upper connected saddles.)



**Figure:**  $x_1$  is a lower connected saddle and  $x_2$  is an upper connected saddle.



## Proposition

Let  $f$  satisfy the basic assumptions. There exists a function  $p_{s^-} : \mathbb{R} \rightarrow [0, \infty)$  such that the following holds. Let  $\Omega \subset \mathbb{R}^2$  and let  $N_{s^-}[\ell, \infty)$  denote the number of lower connected saddles of  $f$  in  $\Omega$  with level above  $\ell$ . Then

$$\mathbb{E}[N_{s^-}[\ell, \infty)] = \text{Area}(\Omega) \int_{\ell}^{\infty} p_{s^-}(x) dx.$$

Analogous statements hold for local maxima, local minima, upper connected saddles and saddles with the densities  $p_{m^+}$ ,  $p_{m^-}$ ,  $p_{s^+}$  and  $p_s$  respectively. These functions can be chosen to satisfy  $p_{s^-} + p_{s^+} = p_s$ , and such that  $p_{m^+}$ ,  $p_{m^-}$  and  $p_s$  are continuous.


## Theorem

Let  $f$  be a Gaussian field satisfying the basic assumptions, and let  $\rho_{m^+}$ ,  $\rho_{m^-}$ ,  $\rho_{s^+}$ ,  $\rho_{s^-}$  denote the critical point densities defined above. Then

$$c_{NS}(\rho, \ell) = \int_{\ell}^{\infty} \rho_{m^+}(x) - \rho_{s^-}(x) + \rho_{s^+}(x) - \rho_{m^-}(x) dx \quad (1)$$

$$c_{ES}(\rho, \ell) = \int_{\ell}^{\infty} \rho_{m^+}(x) - \rho_{s^-}(x) dx \quad (2)$$


and hence  $c_{NS}$  and  $c_{ES}$  are absolutely continuous in  $\ell$ . In addition  $c_{NS}$  and  $c_{ES}$  are jointly continuous in  $(\rho, \ell)$  provided  $\rho$  has a fixed compact support.



**Figure:** On raising the level through the local maximum  $x_1$ , the number of level set components decreases by one. On passing through the local minimum  $x_2$ , the number of level set components increases by one.

# Proof: Intuition

## Lower connected saddle points



**Figure:** On raising the level through the lower connected saddle point  $x_3$ , the number of level set components increases by one.

## Proposition (Cheng-Schwartzman 2017)

Let  $f$  be the random plane wave (RPW) so that  $\kappa(t) = J_0(|t|)$  (the 0-th Bessel function), then

$$\begin{aligned} p_{m^+}(x) &= p_{m^-}(-x) = \frac{1}{4\sqrt{2}\pi^{3/2}} \left( (x^2 - 1)e^{-\frac{x^2}{2}} + e^{-\frac{3x^2}{2}} \right) \mathbb{1}_{x \geq 0} \\ p_s(x) &= \frac{1}{4\sqrt{2}\pi^{3/2}} e^{-\frac{3x^2}{2}}. \end{aligned}$$

Substituting these expressions into the main integral equality and considering the number of 'flip points' (see Kurlberg-Wigman 2018) shows that


## Corollary

Let  $f$  be the RPW and  $\ell \geq 0$ , then


$$\frac{1}{4\pi} \ell \phi(\ell) \leq c_{ES}(\ell) \leq c_{NS}(\ell) \leq \frac{1}{4\pi} \phi(\ell) \left( \sqrt{2} \phi(\sqrt{2}\ell) + \ell \left( 2\Phi(\sqrt{2}\ell) - 1 \right) \right)$$

# Consequences of main results

## Bounds on $c_{NS}$ and $c_{ES}$ in the isotropic case



(a)  $c_{ES}(\rho, \ell)$  for the RPW



(b)  $c_{NS}(\rho, \ell)$  for the RPW


**Figure:** Lower bounds (solid) and upper bounds (dashed) for  $c_{ES}(\rho, \ell)$  and  $c_{NS}(\rho, \ell)$  respectively for the RPW.

The bound on  $c_{ES}(\rho, \ell)$  for  $\ell < 0$  is a result of the equality  $c_{NS}(\rho, \ell) = c_{ES}(\rho, \ell) + c_{ES}(\rho, -\ell)$  and the fact that  $c_{ES}(\rho, \ell)$  is non-decreasing for  $\ell < 0$  (this part is specific to the RPW).


# Consequences of main results

## Bounds on $c_{NS}$ and $c_{ES}$ in the isotropic case

Similar results hold for all isotropic fields satisfying the basic assumptions. (The general expression for upper and lower bounds becomes more complicated, but depends only on the derivatives of  $\kappa$  at 0.)



(a)  $c_{ES}(\rho, \ell)$  for the Bargmann-Fock field.



(b)  $c_{NS}(\rho, \ell)$  for the Bargmann-Fock field.

**Figure:** Lower bounds (solid) and upper bounds (dashed) for  $c_{ES}(\rho, \ell)$  and  $c_{NS}(\rho, \ell)$  respectively, where  $\rho$  is the spectral measure of the Bargmann-Fock field.

### Proposition

Let  $f$  be the Gaussian field with spectral measure

$\rho = \alpha\delta_0 + \frac{\beta}{2}(\delta_K + \delta_{-K}) + \frac{\gamma}{2}(\delta_L + \delta_{-L})$  where  $\beta, \gamma > 0$ ,  $\alpha = 1 - \beta - \gamma \geq 0$  and  $K, L \in \mathbb{R}^2$  are linearly independent. Then

$$c_{NS}(\ell) = |K \times L| \cdot \mathbb{P}(|Y_1 - Y_2| \leq \ell + X_0 \leq Y_1 + Y_2),$$

$$c_{ES}(\ell) = |K \times L| \cdot \mathbb{P}(|Y_1 - Y_2| \leq |\ell + X_0| \leq Y_1 + Y_2),$$

$\times$  denotes the cross product,  $X_0 \sim \mathcal{N}(0, \alpha)$ ,  $Y_1 \sim \text{Ray}(\sqrt{\beta})$ ,  $Y_2 \sim \text{Ray}(\sqrt{\gamma})$  and  $X_0, Y_1, Y_2$  are independent.

If  $c_{NS}(\ell) \neq 0$  then  $N_{LS,R}(\ell)/(\pi R^2)$  converges in  $L^1$  to a non-constant random variable and hence does not converge a.s. to a constant, and this statement also holds for  $c_{ES}$  and  $N_{ES,R}(\ell)/(\pi R^2)$ . Furthermore

$$p_{m^+}(x) = p_{m^-}(-x) = |K \times L| \cdot p_{X_0+Y_1+Y_2}(x)$$


$$p_{s^-}(x) = p_{s^+}(-x) = |K \times L| \cdot p_{X_0+|Y_1-Y_2|}(x)$$

where  $p_Z$  denotes the probability density of a random variable  $Z$ .



# Consequences of main results


## Derivation of $c_{NS}$ and $c_{ES}$ for 4 point spectral measures



**Figure:** The functions  $c_{ES}(\ell)$  (left) and  $c_{NS}(\ell)$  (right) with  $\alpha = 0$  for  $\beta - \gamma = 0$  (solid),  $\beta - \gamma = 0.5$  (dashed) and  $\beta - \gamma = 0.9$  (dotted) respectively.

# Consequences of main results

## Derivation of $c_{NS}$ and $c_{ES}$ for 5 point spectral measures



**Figure:** The functions  $c_{ES}(\ell)$  (left) and  $c_{NS}(\ell)$  (right) with  $\beta = \gamma$  for  $\alpha = 0.1$  (solid),  $\alpha = 0.3$  (dashed) and  $\alpha = 0.6$  (dotted) respectively.

1. Characterising  $p_{s-}$  (or  $p_{s+}$ )
2. Higher dimensions
3. Continuous differentiability of  $c_{NS}$
4. Bimodality

