The Wiener chaos expansion for non-local functionals of Gaussian fields

Michael McAuley *Technological University Dublin* Based on joint work with Stephen Muirhead

> Conference PRIN GRAFIA, Cortona, 19th June 2025

Slides available at https://michael-mcauley.github.io



Outline

1. Introduction

2. Wiener chaos method for local functionals

3. Wiener chaos method for non-local functionals



Smooth Gaussian fields

- Let $f : \mathbb{R}^d \to \mathbb{R}$ be a stationary C^2 Gaussian field with mean zero and variance one.
- The distribution of f is specified by its covariance function K : ℝ^d → ℝ defined as

$$K(x-y) = \operatorname{Cov}[f(x), f(y)] \qquad \forall x, y \in \mathbb{R}^d.$$

We will consider the geometry/topology of the excursion sets

$$\{f \ge \ell\} := \left\{ x \in \mathbb{R}^d \mid f(x) \ge \ell \right\}$$
 for $\ell \in \mathbb{R}$.



Figure: Excursion sets $\{f \ge 0\}$ in white for the fields on \mathbb{R}^2 with $K(x) = J_0(|x|)$, the 0-th Bessel function, (left) and $K(x) = \exp(-|x|^2/2)$ (right).



Motivation: Percolation theory

- Percolation theory studies the large scale topological properties of spatial random models.
- ▶ Phase transition: for a given field, there is a critical level *l_c* such that, with probability one
 - for ℓ > ℓ_c, {f ≥ ℓ} contains only bounded components,
 - for $\ell < \ell_c$, $\{f \ge \ell\}$ contains a unique unbounded component.

See [1] for a survey.



Figure: The excursion sets $\{f \ge \ell\}$ for $\ell = 0.05$ (left), $\ell = 0$ (middle) and $\ell = -0.05$ (right). Largest component highlighted in green.



Local vs non-local functionals

A functional of a random field is described as **local** if it is an integral of a pointwise function of the field and its derivatives:

$$\int_D \varphi(f(x), \nabla f(x), \nabla^2 f(x)) \ \mu(dx)$$

Examples

Local functionals

- volume $\int_D \mathbb{1}_{f(x) \ge \ell} dx$
- boundary volume $\int_D \mathbb{1}_{f(x)=\ell} \mathcal{H}^{d-1}(dx)$
- Euler characteristic

Non-local functionals

- number of connected components
- Betti numbers
- Volume of the unbounded component
- We focus on non-local functionals which are 'approximately additive' over the domain.



Local vs non-local functionals

What is known?

Local functionals

- Powerful methods available including the Kac-Rice formula and the Wiener chaos expansion,
- Can typically characterise the mean, variance and asymptotic distribution,
- Results known for a wide variety of covariance structures.

Non-local functionals

▶ No unifying theory, but many partial results using a variety of methods:

Type of result	Methods
Law of large numbers [10]	Ergodic argument
Variance bounds[11, 4, 3]	Coupling, interpolation formulae
Central limit theorem [2, 8, 7]	Martingale techniques

Most results are sub-optimal or hold only for a restricted class of covariance functions.



Outline

1. Introduction

2. Wiener chaos method for local functionals

3. Wiener chaos method for non-local functionals



A classical problem

Consider a functional of the form

$$F_R = \int_{[-R,R]^d} \varphi(f(x) - \ell) \, dx$$

for some $\varphi : \mathbb{R} \to \mathbb{R}$.

▶ For simplicity, assume that

$$\mathcal{K}(x)\sim c|x|^{-lpha}$$
 as $|x|
ightarrow\infty$

for some $\alpha \in (0, d)$.

Question: Can we describe the asymptotic statistics (mean, variance, distribution) of F_R as $R \to \infty$?

By Fubini's theorem,

$$\mathbb{E}[F_R] = (2R)^d \mu(\ell)$$

where $\mu(\ell) := \mathbb{E}[\varphi(f(0) - \ell)].$



Wiener chaos expansion

Let \mathcal{G} be a set of centred jointly Gaussian variables. Let \mathcal{P}_n be the space of all polynomials of degree $\leq n$ in \mathcal{G} . The *n*-th Wiener chaos of \mathcal{G} is $\mathcal{G}^{:n:} := \overline{\mathcal{P}_n} \cap \overline{\mathcal{P}_{n-1}}^{\perp}$.

Theorem

Let the random variable F be square integrable and $\sigma(\mathcal{G})$ -measurable, then

$$F \stackrel{L^2}{=} \sum_{n=0}^{\infty} Q_n[F]$$

where Q_n denotes projection onto $\mathcal{G}^{:n:}$.



Hermite polynomials

▶ The Hermite polynomials $(H_n)_{n\geq 0}$ can be defined inductively by setting

$$H_0(x) = 1$$
 and $H_{n+1}(x) = xH_n(x) - H'_n(x)$.

Properties:

1. If X, Y are jointly normal with mean zero and variance one then

$$\mathbb{E}[H_n(X)H_m(Y)] = \begin{cases} n! \operatorname{Cov}[X, Y]^n & \text{if } n = m \\ 0 & \text{if } n \neq m. \end{cases}$$

Hence if $X \in \mathcal{G}$ then $H_n(X) \in \mathcal{G}^{:n:}$. 2. If $\mathbb{E}[\varphi^2(Z)] < \infty$ for $Z \sim \mathcal{N}(0, 1)$ then

$$\varphi = \sum_{n=0}^{\infty} a_n H_n$$

where $\sum_{n} a_n^2 n! < \infty$.



Chaos expansion for a local functional Variance asymptotics

• Considering the expansion $\varphi(\cdot - \ell) = \sum_n a_n(\ell) H_n$ yields

$$F_{R} = \sum_{n=0}^{\infty} a_{n}(\ell) \int_{[-R,R]^{d}} H_{n}(f(x)) \ dx = \sum_{n=0}^{\infty} Q_{n}[F_{R}].$$

• The variance of F_R can be computed by considering

$$\operatorname{Var}[Q_n[F_R]] = a_n(\ell)^2 \iint_{[-R,R]^{2d}} \operatorname{Cov}[H_n(f(x)), H_n(f(y))] \, dxdy$$
$$= a_n(\ell)^2 n! \iint_{[-R,R]^{2d}} K(x-y)^n \, dxdy$$

for $n \ge 1$. Since $K(x) \sim c|x|^{-\alpha}$, for $n \ge 1$

$$\operatorname{Var}[Q_n[F_R]] \sim a_n(\ell)^2 c_{K,n} \times \begin{cases} R^{2d-n\alpha} & \text{if } n\alpha < d, \\ R^d \log R & \text{if } n\alpha = d, \\ R^d & \text{if } n\alpha > d. \end{cases}$$



Chaos expansion for a local functional

Convergence in distribution

• Since
$$H_1(x) = x$$

 $Q_1[F_R] = a_1(\ell) \int_{[-R,R]^d} f(x) dx$

which is Gaussian.

• For $n\alpha \ge d$, one can show that

$$\frac{Q_n[F_R]}{\sqrt{\operatorname{Var}[Q_n[F_R]]}} \xrightarrow{d} \mathcal{N}(0,1)$$

using the **fourth-moment theorem** (or method of moments) and a **diagram formula** for the moments of Hermite polynomials.

For 1 < n < d/α by expressing Hermite polynomials as multiple Wiener-Itô integrals one can show

$$\frac{Q_n[F_R]}{\sqrt{\operatorname{Var}[Q_n[F_R]]}} \xrightarrow{d} c_{K,n} \int \mathcal{F}[\mathbbm{1}_{[-1,1]^d}](\sum_{i=1}^n u_i) \prod_{i=1}^n \frac{W(du_i)}{|u_i|^{(d-\alpha)/2}}$$

where the latter follows a Hermite distribution.



Conclusion: Limit theorems for a local functional

Theorem (Breuer-Major/Dobrushin-Major theorem) Let $n^*(\ell) = \inf\{n : a_n(\ell) \neq 0\}$. If f satisfies some technical conditions, then

$$\operatorname{Var}[F_R] \sim c_{\mathcal{K},\varphi,\ell} \times \begin{cases} R^{2d-n^*\alpha} & \text{if } n^*\alpha < d, \\ R^d \log R & \text{if } n^*\alpha = d, \\ R^d & \text{if } n^*\alpha > d, \end{cases} \quad \text{as } R \to \infty.$$

Moreover if $n^* = 1$ or $n^* \alpha \ge d$ then

$$\frac{F_R - \mu(\ell)}{\sqrt{\operatorname{Var}[F_R]}} \stackrel{d}{\to} \mathcal{N}(0, 1).$$

For other values of n^* , the limiting distribution is a Hermite distribution.

Remark

- Typically n^{*}(ℓ) = 1 for all but finitely many values of ℓ, which are described as anomalous levels.
- If φ is regular then a_n(ℓ) = (−1)ⁿμ⁽ⁿ⁾(ℓ)/n! so that anomalous levels correspond to critical points of μ.



Outline

1. Introduction

2. Wiener chaos method for local functionals

3. Wiener chaos method for non-local functionals



Wiener chaos method for non-local functionals $_{\mathsf{Setting}}$

In joint work with Stephen Muirhead [9], we adapt the Wiener chaos method to prove limit theorems for a non-local functional.

▶ Let $f : \mathbb{Z}^d \to \mathbb{R}$ be the Gaussian free field (in $d \ge 3$), so that

$$K(x-y) \sim c_d |x-y|^{-(d-2)}.$$

The cluster count N_R(f) is the number of clusters (i.e. connected components) of the graph {f ≥ ℓ} ∩ [−R, R]^d.



Part 1: Identifying chaos projections Smooth functionals

Proposition

Let $D \subset \mathbb{Z}^d$ be finite and $\Phi \in C^{\infty}(\mathbb{R}^D)$, then

$$Q_n[\Phi(f)] = \frac{1}{n!} \sum_{x_1,\ldots,x_n \in D} \mathbb{E}[\partial_{x_1} \ldots \partial_{x_n} \Phi(f)] : f(x_1) \ldots f(x_n):$$

where the Wick polynomial $:f(x_1) \dots f(x_n)$: is defined as $Q_n[f(x_1) \dots f(x_n)]$.

- Proof 1: Elementary argument using Gaussian integration by parts.
- Proof 2: Stroock formula:

$$Q_n[\Phi(f)] = I_n\left(\frac{1}{n!}\mathbb{E}[D^n\Phi(f)]\right)$$

= $\frac{1}{n!}\sum_{x_1,\ldots,x_n\in D}\mathbb{E}[\partial_{x_1}\ldots\partial_{x_n}\Phi(f)]I_n(e_1\otimes\cdots\otimes e_n)$



Part 1: Identifying chaos projections Cluster count

The **discrete derivative** d_x is defined as

$$d_x N_R(f) = N_R(\{f \ge \ell\} \cup \{x\}\}) - N_R(\{f \ge \ell\} \setminus \{x\}).$$

Let $\Lambda_R = [-R, R]^d \cap \mathbb{Z}^d$.

Proposition

For $R \ge 1$

$$Q_n[N_R(f)] = \frac{1}{n!} \sum_{x_1,\ldots,x_n \in \Lambda_R} P_R(x_1,\ldots,x_n) : f(x_1)\ldots f(x_n):$$

where the **pivotal intensity** P_R is defined for distinct points $\underline{x} = (x_1, \ldots, x_n)$ as

$$P_R(\underline{x}) = \mathbb{E}[d_{x_1} \dots d_{x_n} N_R(f) | f(\underline{x}) = \ell] \varphi_{f(\underline{x})}(\ell),$$

and $\varphi_{f(\underline{x})}$ is the density of $f(\underline{x})$.



Part 1: Identifying chaos projections Part 2: Semi-locality of pivotal intensities

Comparing with a local functional:

Local: $Q_n[F_R] = a_n(\ell) \sum_{x \in \Lambda_R} : f(x)^n:$

Non-local: $Q_n[N_R(f)] = \frac{1}{n!} \sum_{x_1, \dots, x_n \in \Lambda_R} P_R(x_1, \dots, x_n) : f(x_1) \dots f(x_n):$

since $:f(x)^{n}: = H_{n}(f(x)).$

- ▶ Hence the local case corresponds to $P_R(x_1, ..., x_n) = n!a_n(\ell)\mathbb{1}_{x_1=\cdots=x_n}$.
- One could imagine extending the analysis from the local case if P_R is approximately stationary and has rapid off-diagonal decay.
- We refer to this as semi-locality.



Part 1: Identifying chaos projections Part 2: Semi-locality of pivotal intensities

• Recall that for distinct points $\underline{x} = (x_1, \dots, x_n)$

$$P_R(\underline{x}) = \mathbb{E}[d_{x_1} \dots d_{x_n} N_R(f) | f(\underline{x}) = \ell] \varphi_{f(\underline{x})}(\ell).$$



Figure: For this configuration $d_{x_1}d_{x_2}N_R(f) = 1$ but $d_{x_1}d_{x_3}N_R(f) = 0.$

In general, if d_{x1}...d_{xn}N_R(f) ≠ 0 then x1,...,xn must be joined by bounded clusters of {f ≥ ℓ}.



Part 1: Identifying chaos projections Part 2: Semi-locality of pivotal intensities

Theorem (Truncated arm decay [6])

Let $f : \mathbb{Z}^d \to \mathbb{R}$ be the Gaussian free field for $d \ge 3$. There exists $\ell_c \in \mathbb{R}$ such that for every $\ell \neq \ell_c$, the probability that 0 is contained in a bounded cluster of $\{f \ge \ell\}$ of diameter at least n is at most $e^{-cn^{\rho}}$ for some $c, \rho > 0$.

Corollary

For $\ell \neq \ell_c$ there exists $c, C, \rho > 0$ such that

$$P_R(\underline{x}) \leq C e^{-c \operatorname{diam}(\underline{x})^{
ho}}$$

where $diam(\underline{x})$ denotes the diameter of \underline{x} .



Part 1: Identifying chaos projections

Part 2: Semi-locality of pivotal intensities

Part 3: Convergence of semi-local chaoses

- Arguments for local functionals (based on the fourth-moment theorem/multiple Wiener-Itô integrals) can be extended to the semi-local case.
- Calculations involving covariance kernels become more involved but are conceptually straightforward:

Local:
$$\operatorname{Var}[Q_n[F_R]] = a_n(\ell)^2 \sum_{x,y \in \Lambda_R} K(x-y)^n$$

Non-local:
$$\operatorname{Var}[Q_n[N_R(f)]] = \frac{1}{(n!)^2} \sum_{\underline{x}, \underline{y} \in \Lambda_R^n} P_R(\underline{x}) P_R(\underline{y}) \prod_{i=1}^n K(x_i - y_i).$$

► To control the tail of the chaos expansion we use an **interpolation formula** for $Var[\sum_{n \ge N} Q_n[N_R(f)]]$ in terms of discrete derivatives of order N.



Conclusion: Limit theorems for the cluster count

We define the mean clusters-per-vertex as

$$\mu(\ell) := \lim_{R \to \infty} \frac{\mathbb{E}[N_R(f)]}{(2R)^d}.$$

Theorem

Let $f:\mathbb{Z}^3\to\mathbb{R}$ be the Gaussian free field and $\ell\neq\ell_c.$

$$\operatorname{Var}[N_{R}(f)] \sim c_{\ell} \times \begin{cases} R^{5} & \text{if } \mu'(\ell) \neq 0 \\ R^{4} & \text{if } \mu'(\ell) = 0, \mu''(\ell) \neq 0 \\ R^{3} \log R & \text{if } \mu'(\ell) = \mu''(\ell) = 0, \mu'''(\ell) \neq 0 \\ R^{3} & \text{otherwise.} \end{cases}$$

In the second case, the (normalised) limiting distribution is a Hermite distribution, in all other cases it is Gaussian.

- Analogous results hold for d ≥ 4 and other fields but are omitted here for brevity.
- Similar to results in local case, but the requirement that $\ell \neq \ell_c$ is new. **T**

Summary

Open questions:

- Can this approach be extended to smooth fields?
- Does this approach enable the Malliavin-Stein method for non-local functionals?
- What happens at the critical level?

Thank you for listening!



Bibliography I

- D. Beliaev. "Smooth Gaussian fields and percolation". In: Probability Surveys (2023). URL: https://doi.org/10.1214/23-PS24.
- [2] D. Beliaev, M. McAuley, and S. Muirhead. "A central limit theorem for the number of excursion set components of Gaussian fields". In: Ann. Probab. (2024). URL: https://doi.org/10.1214/23-A0P1672.
- D. Beliaev, M. McAuley, and S. Muirhead. "A covariance formula for the number of excursion set components of Gaussian fields and applications". In: Ann. Inst. Henri Poincaré Probab. Stat. (2025). URL: https://doi.org/10.1214/23-aihp1430.
- [4] D. Beliaev, M. McAuley, and S. Muirhead. "Fluctuations of the number of excursion sets of planar Gaussian fields". In: *Probab. Math. Phys.* (2022). URL: https://doi.org/10.2140/pmp.2022.3.105.
- [5] D. Beliaev, S. Muirhead, and A. Rivera. "A covariance formula for topological events of smooth Gaussian fields". In: Ann. Probab. (2020). URL: https://doi.org/10.1214/20-A0P1438.
- [6] H. Duminil-Copin et al. "Equality of critical parameters for percolation of Gaussian free field level sets". In: Duke Math. J. (2023). URL: https://doi.org/10.1215/00127094-2022-0017.

Bibliography II

- [7] C. Hirsch and R. Lachièze-Rey. "Functional central limit theorem for topological functionals of Gaussian critical points". In: arXiv preprint arXiv:2411.11429 (2024).
- [8] M. McAuley. "Three central limit theorems for the unbounded excursion component of a Gaussian field". In: arXiv preprint arXiv:2403.03033 (2024).
- [9] M. McAuley and S. Muirhead. Limit theorems for the number of sign and level-set clusters of the Gaussian free field. 2025. URL: https://arxiv.org/abs/2501.14707.
- [10] F. Nazarov and M. Sodin. "Asymptotic Laws for the Spatial Distribution and the Number of Connected Components of Zero Sets of Gaussian Random Functions". In: Journal of Mathematical Physics, Analysis, Geometry (2016). URL: https: //jmag.ilt.kharkiv.ua/index.php/jmag/article/view/jm12-0205e
- [11] F. Nazarov and M. Sodin. "Fluctuations in the number of nodal domains". In: J. Math. Phys. (2020). URL: https://doi.org/10.1063/5.0018588.